



Allergy & Dermatology

s p e c i a l i s t s

Mohs Post Operative Wound Care

- Leave the bandage on throughout the night.
- Remove the bandage the next day.
- Wash the wound daily with clean soap and water only, no peroxide. Showers are fine. Apply a thin layer of either Vaseline or Polysporin to the area twice daily until the next appointment.
- Do not use Neosporin!
- A little bleeding or seeping from the wound is normal. If there is any significant bleeding noticed, apply direct pressure to the area for 20 minutes. You may place an ice pack on the area for additive pressure to help the blood clot. If bleeding persists call the numbers available below or go to the nearest emergency room.
- A small amount of swelling, redness, and tenderness is to be expected. If you experience worsening pain, swelling or redness please call the office for a follow up.
- A bag is being provided with all the necessary supplies needed to last for at least a couple of days or up until your next appointment. Use the Aquaphor/Vaseline on the wound, cut up the Telfa pads and Hypafix tape to size, and use the ice pack as needed for pain and/or swelling. You can also apply a band-aid or other dressings if appropriate.
- If you have any questions, comments, or concerns please don't hesitate to call the numbers below:

Phoenix - (602) 277-7686

Sun City West - (623) 544-7755

Sun City - (623) 977-4218

Arrowhead - (623) 376-7600

- If you are having problems after hours, our answering service will be able to contact Dr. Juacek and have him call you back shortly. They can be reached at (480) 980-3887.